

## Biliary Drain/Stent Insertion

You have received the following Anaesthetic that is checked ( ✓ )

- Local Anesthetic** – An injection of anesthesia (freezing) to numb a small area at the site of your biopsy/procedure. It is common to feel mild or temporary numbness or tingling around the site.
- IV Sedation** - Medications given through your intravenous to relax you and decrease anxiety while you are still awake. It is common to have trouble remembering what happened during your procedure, and/or feel tired for rest of the day. You must return home with a responsible adult who will remain with you overnight. You should not take public transit after the procedure. **With IV sedation, do not drive, use machinery or power tools or make important legal or financial decisions for the next 24 hours.**

### Diet

There are no diet restrictions. Avoid alcohol for 24 hours. Drink plenty of fluids for the next 24 hours.

### Activity

Get plenty of rest for 24 hours after the procedure. Avoid any strenuous exercise, lifting objects weighing more than 5 pounds for 7 days, or **activity that may dislodge the drain.**

### Dressing

Keep your bandage dry. Replace the bandage as needed if it becomes soiled or wet. If you have a drainage bag empty as required. You may bathe or shower the day after your procedure, ensuring the drain site is covered and remains dry. **Do NOT swim or use a hot tub while you have a drain inserted.**

### Medication

Avoid taking medication containing aspirin or blood thinners for 24 hours after your procedure unless otherwise directed. Take pain medication, as prescribed, or you may take Acetaminophen (Tylenol®) (if no allergy) 1 – 2 tablets every 4 – 6 hours as needed for pain, up to a maximum of 4,000 mg in 24 hours. You may take your next pain medication at: \_\_\_\_\_.

### Follow Up

If applicable Home and Community services will contact you to set up home care services, call 905-762-4028 if you do not hear from them in 24 hours.

**CALL YOUR DOCTOR DURING OFFICE HOURS OR GO TO THE NEAREST EMERGENCY DEPT. IF:**

1. Fever over 38.5°C or 101.3°F, or shakes and chills
2. Increasing redness, tenderness, swelling, bleeding or bruising at the drain site
3. Increasing pain to the abdomen not relieved by acetaminophen or your prescribed pain medication, or increased abdominal distension
4. Bright red blood draining into the drainage bag (“pink tinged” fluid is ok)
5. Chest pain, shortness of breath, difficulty breathing, or heart rate over 100 bpm
6. Dizziness, light headedness or confusion, exhaustion, extreme weakness, sweating, pale/blue skin
7. Persistent nausea or vomiting

**Diagnostic Imaging Day Unit 905-472-7373 ext. 1521 (Mon-Fri 8:00 am-3:00 pm)**