

Port-A-Cath Removal

You have received the following Anaesthetic that is checked (✓)

- Local Anesthetic** - An injection of anesthesia (freezing) to numb a small area at the site of your procedure. It is common to feel mild or temporary numbness or tingling around the site.
- IV Sedation** - Medications given through your intravenous to relax you and decrease anxiety while you are still awake. It is common to have trouble remembering what happened during your procedure, and/or feel tired for the rest of the day. You must return home with a responsible adult who will remain with you overnight. You should not take public transit after the procedure.

With IV sedation, do not drive, use machinery or power tools or make important legal or financial decisions for the next 24 hours.

Diet

There are no diet restrictions. Avoid alcohol for 24 hours. Drink plenty of fluids for the next 24 hours.

Activity

Get plenty of rest for 24 hours after the procedure, then return to usual activities (school, work and home) as you feel able.

Dressing

Keep your bandage dry. You may remove the outside bandage after 72 hours. You may have paper tapes over your incision – leave these on until they fall off on their own. The stitches are usually dissolvable – you will be instructed if there are stitches to remove. Do not wear a tight bra or suspenders. You may bathe or shower the day after your procedure, ensuring the bandages are covered and remain dry. Do not immerse the incision in water until it is fully healed. **Do NOT swim or use a hot tub or immerse the incision in water until it is fully healed.**

Medication

Avoid taking medication containing aspirin or blood thinners for 24 hours after your procedure unless otherwise directed. Take pain medication, as prescribed, or you may take Acetaminophen (Tylenol®) (if no allergy) 1 – 2 tablets every 4 – 6 hours as needed for pain, up to a maximum of 4,000 mg in 24 hours. You may take your next pain medication at: _____.

Follow Up

Return to see your Oncologist or Doctor as directed.

CALL YOUR DOCTOR DURING OFFICE HOURS, OR GO TO THE NEAREST EMERGENCY DEPARTMENT OR CALL 911 IF YOU HAVE:

1. Fever over 38.5°C or 101.3°F, or shakes and chills
2. Increasing pain, redness, swelling, discharge or bleeding around your incision, or swelling to the neck or face on the side of incision
3. Chest pain, shortness of breath, difficulty breathing, or heart rate over 100 bpm
4. Dizziness, light headedness or confusion, exhaustion, extreme weakness, sweating, pale/blue skin
5. Persistent nausea or vomiting.

Diagnostic Imaging Day Unit 905-472-7373 ext. 1521 (Mon-Fri 8:00 am-3:00 pm)