

Interventional Radiology

Discharge Instructions

Tunneled Abdominal Drain Removal

You have received the following Anaesthetic that is checked (✓)

- Local Anesthetic** – An injection of anesthesia (freezing) to numb a small area at the site of your biopsy/procedure. It is common to feel mild or temporary numbness or tingling around the site.
- IV Sedation** - Medications given through your intravenous to relax you and decrease anxiety while you are still awake. It is common to have trouble remembering what happened during your procedure, and/or feel tired for rest of the day. You must return home with a responsible adult who will remain with you overnight. You should not take public transit after the procedure. **With IV sedation, do not drive, use machinery or power tools or make important legal or financial decisions for the next 24 hours.**

Diet

There are no diet restrictions. Avoid alcohol for 24 hours.

Activity

Get plenty of rest for 24 hours after the procedure. Return to usual activities (school, work and home) as you feel able.

Dressing

Keep your bandage dry. Replace bandage as needed if it becomes soiled or wet. Bandage may be removed after 48 hours if no oozing. You may bathe or shower the day after your procedure, ensuring the bandage is covered and remains dry. **Do NOT swim or use a Hot tub until drain site is healed.**

Medication

Avoid taking medication containing aspirin or blood thinners for 24 hours after your procedure unless otherwise directed. Take pain medication, as prescribed, or you may take Acetaminophen (Tylenol®) (if no allergy) 1 – 2 tablets every 4 – 6 hours as needed for pain, up to a maximum of 4,000 mg in 24 hours. You may take your next pain medication at: _____.

Follow Up

Return to see your doctor as directed. Home and Community services will continue their services as needed or as ordered by your doctor.

CALL YOUR DOCTOR DURING OFFICE HOURS OR GO TO THE NEAREST EMERGENCY DEPT. IF:

1. Fever over 38.5°C or 101.3°F, or shakes and chills
2. Increasing pain, redness, swelling, bleeding or bruising at the procedure site
3. Increasing pain to the abdomen not relieved by acetaminophen or abdominal distention
4. Chest pain, shortness of breath, difficulty breathing, or heart rate over 100 bpm
5. Dizziness, light headedness or confusion, exhaustion, extreme weakness
6. Persistent nausea or vomiting

Diagnostic Imaging Day Unit 905-472-7373 ext. 1521 (Mon-Fri 8:00 am-3:00 pm)